

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Powdery mildew caused by *Golovinomyces cichoracearum* on eastern purple coneflower (*Echinacea purpurea*) in Italy**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1680218> since 2019-03-29T18:41:22Z

*Published version:*

DOI:10.1094/PDIS-10-17-1691-PDN

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**Powdery Mildew caused by *Golovinomyces cichoracearum* on Eastern Purple Coneflower (*Echinacea purpurea*) in Italy.**

A. Garibaldi, D. Bertetti, and S. Matić, Centre of Competence AGROINNOVA, University of Torino, Largo Braccini 2, 10095 Grugliasco, Italy; and M. L. Gullino, Centre of Competence AGROINNOVA and DISAFA, University of Torino, Largo Braccini 2, 10095 Grugliasco, Italy.

Eastern purple coneflower (*Echinacea purpurea*) is a herbaceous plant grown as ornamental and for its medicinal properties. During the summer 2017 and the following autumn, 6-8-months-old potted plants of *E. purpurea* grown in a private garden located in Biella province (northern Italy, showed symptoms and signs of a previously unknown powdery mildew. A white mycelium covered stems, leaves and inflorescences of all plants. The ligules were particularly affected and dried. Mycelium produced erect conidiophores, with a cylindrical foot cell measuring  $81\text{--}129 \times 11\text{--}13$  (mean  $111 \times 12$ )  $\mu\text{m}$ , followed by 3 shorter cells measuring  $13\text{--}37 \times 10\text{--}17$  (mean:  $24 \times 13$ )  $\mu\text{m}$ . Elliptical conidia were collected in short chains and measured  $30\text{--}42 \times 16\text{--}24$  (mean:  $36 \times 21$ )  $\mu\text{m}$ , length/width (l/w) ratio between 1.3 to 2.6 (mostly 1.5 - 2.0). Fibrosin bodies were absent. The teleomorph was not observed. DNA was extracted from mycelium, conidiophores and conidia collected from leaves and ligules of affected plants using the E.Z.N.A. Fungal DNA Mini Kit (Omega Bio-Tek, Darmstadt, Germany). A PCR reaction was performed using primers ITS1 (White et al. 1990) and PM6 (Takamatsu and Kano 2001), to amplify the Internal Transcribed Spacer (ITS) region of rDNA. Blastn analysis of 498 bp sequence (Gene Bank accession number MG182430) showed a 99% similarity with *Golovinomyces cichoracearum*, *G. spadiceus* and *G. ambrosiae*. Because conidium size observed on *E. purpurea* (in particular: width  $> 20 \mu\text{m}$  and l/w ratio) are more similar to *G. cichoracearum* (Braun and Cook 2012), the causal agent of powdery mildew detected on this host was identified as *G. cichoracearum*. To confirm the pathogenicity, three 8-months-old plants of *E. purpurea* were artificially inoculated by gently pressing mycelium, conidiophores and conidia of affected leaves onto healthy leaves. Three non-inoculated plants served as controls. All the plants were kept at temperatures ranging between 15 and 25°C. About 15 days after the artificial inoculation, the first symptoms of powdery mildew appeared on inoculated leaves on which morphological characteristics of the pathogen were identical to those reported above. Any symptoms appeared on controls. *Erysiphe* (Syn.: *Golovinomyces*) *cichoracearum* was reported on *E. purpurea* in Canada (Sholberg and Ginns 1999) and in Poland. To our knowledge, this is the first report of *G. cichoracearum* on *E. purpurea* in Italy. The economic importance of the disease described on *E. purpurea* is at present limited, but it could spread to several hosts belonging to the Compositae on which *G. cichoracearum* has been reported (Braun and Cook 2012).

## *References*

- Braun, U., and Cook, R. T. A. 2012. Taxonomic Manual of the *Erysiphales* (Powdery Mildews). CBS Biodiversity Series No. 11. CBS Utrecht, The Netherlands.
- Sholberg, P. L., Ginns, J. H., and Li, T. S. C. 1999. Plant Disease 83:694.
- Takamatsu, S., and Kano, Y. 2001. Mycoscience 42:135-139.
- White, T. J., et al. 1990. Page 315. In: PCR Protocols: A Guide to Methods and Applications. M. A. Innis et al., eds. Academic Press Inc., New York.